

University of Groningen

A study of planetary nebulae. Distances and physical properties.

Gathier, Roelof

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1984

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Gathier, R. (1984). *A study of planetary nebulae. Distances and physical properties*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Taylor, K.: 1984, preprint
 Atmospheres and Envelopes, ed. B.
 ger, Berlin, p. 229
 Red Giants, ed. I. Iben Jr., A.
 . Flower, Reidel, Dordrecht, p.
 D.: 1982, Monthly Notices Roy.
 Astron. Soc. J. 274, L61
 19, 108
 103, 119
 708
 Physical Processes in Red Giants, ed.
 Reidel, p. 463
 IAU Symp. 103, ed. D.R. Flower,
 . 103, ed. D.R. Flower, Reidel,
 Astrophys. 130, 269
 Rev. Mex. Astron. Astrof. 2, 181
 Astrophys. J. 234, 217
 , L27
 Astrophys. 121, 77
 ss on Stellar Evolution. ed. C.
 p. 353

CHAPTER 6 SUMMARY

This investigation had two main purposes. The first was to determine individual distances of a number of planetary nebulae (PN) in order to considerably enlarge the sample of PN with accurate distances. The second purpose was to use these distances to investigate several properties of PN such as their intrinsic physical parameters, their evolution in the Hertzsprung-Russell diagram and to verify the validity of various statistical distance scales of PN.

In Chapter 2 a frequently used statistical method for the determination of PN distances, the "Shklovsky method", has been investigated. This study was based on high resolution radio observations of about 40 PN at the galactic centre (Chapter 2A) and of K 648 in the globular cluster M15 (Chapter 2B).

In Chapters 3 and 4 two methods to determine distances to PN have been described and applied. In Chapter 3 distances to 12 PN have been determined from relations of interstellar reddening as a function of distance along the lines of sight towards the PN. In Chapter 4 distances to 11 PN have been derived using 21 cm HI absorption measurements.

In Chapter 5 the distances derived in Chapters 3 and 4 and other accurate distances already published have been combined to a sample of 30 PN with "standard distances". This sample has been used to determine intrinsic parameters of the PN, to study their evolution and to investigate several statistical distance scales.

The main results and conclusions of the studies presented in Chapters 2 to 5 are summarized below.

- 1) Using high resolution 6 cm observations obtained with the VLA, accurate flux densities and angular sizes could be determined for about 40 PN at the galactic centre and for K 648 in M15. The ionized nebular masses derived vary from $\leq 0.008 M_{\odot}$ to $\sim 0.6 M_{\odot}$. These results show that the Shklovsky method for determining distances to PN will generally lead to unreliable distances.
- 2) Accurate Walraven VBLUV photometry of stars in small fields around 13 PN has been obtained. Using theoretical model atmospheres the effective temperature T_{eff} and surface gravity $\log g$ could be derived for program stars with $T_{\text{eff}} \gtrsim 7000$ K. Distances to these stars were determined using a calibration of absolute magnitude as a function of T_{eff} and $\log g$. This calibration has been derived from theoretical evolutionary models. It could be shown that the calibration used leads to quite accurate

distances of the program stars; the systematic errors are probably smaller than $\sim 10\%$. The scatter in the resulting reddening-distance relations along the lines of sight towards the PN could in all but two cases be explained by errors in the distances of individual program stars. This shows that the irregular distribution of dust across the sky is not necessarily the limiting factor in the application of the reddening-distance method, if stars within a few tenths of a degree from the PN are used.

The values for the interstellar extinction in the direction of the PN studied have been determined accurately. Several methods were used such as a comparison of the H β flux with the radio continuum flux density and the strength of the absorption feature at λ 2200 Å in the UV spectra of PN, obtained with the IUE satellite. Using infrared observations obtained by IRAS, it has been shown that internal reddening is generally negligible compared to the foreground reddening of PN.

The reddening-distance relations, combined with the reddenings of the PN, provided distances to 12 PN with accuracies ranging from ~ 10 to $\sim 40\%$.

- 3) HI absorption measurements of 24 PN together with absorption measurements of background sources near the lines of sight towards the PN have been obtained using the Westerbork radio telescope. HI emission spectra in the directions to the PN have been obtained with the Dwingeloo radio telescope. Using the kinematic distance information derived from these measurements, combined with distance information from other sources, it was possible to determine distances to 11 PN with accuracies ranging from ~ 25 to $\sim 40\%$.
- 4) The distances derived with the reddening-distance method and by means of the HI absorption measurements, combined with distances already published, a sample of 30 "standard distances" has been constructed. These distances are accurate within $\sim 40\%$. At present this is the largest sample of accurate distances to PN that is available.
- 5) Using the sample of standard distances it was possible to give accurate positions of the PN in the Hertzsprung-Russell diagram. A comparison of these positions with theoretical evolutionary tracks shows that there are severe discrepancies between theory and observations. The observations show a considerable spread in central star luminosities while theoretical models predict a narrow range in luminosity. It is shown that the most plausible explanation for this discrepancy is that the evolutionary time scales predicted by theory are not correct. It is suggested that these time scales can be adjusted by using different theoretical values for M_{eN} , the remnant envelope mass of the central star at $T_{\text{eff}} = 3 \cdot 10^4$ K. Adjustment of M_{eN} can be accomplished by a

revision of
ejection, a
superwind p

The obs
imply that
found earli
tribution o
tional fact
parable.

The theo
high-luminos
firmed by th

6) It is found
ionized mas
results of th

7) The standard
tistical dist
from these s
large spread
central star
all PN canno
mixture of s
stars with di

ors are probably small-
ddening-distance rela-
d in all but two cases
individual program stars.
t across the sky is not
ion of the reddening-
a degree from the PN

n the direction of the
eral methods were used
io continuum flux den-
at λ 2200 Å in the UV
Using infrared observa-
internal reddening is
ddening of PN.

With the reddenings of
es ranging from ~ 10 to

th absorption measure-
ght towards the PN have
pe. HI emission spectra
with the Dwingeloo radio
ion derived from these
from other sources, it
ith accuracies ranging

method and by means of
tances already publish-
en constructed. These
t this is the largest
le.

sible to give accurate
agram. A comparison of
acks shows that there
vations. The observa-
ar luminosities while
nosity. It is shown
crepancy is that the
e not correct. It is
d by using different
e mass of the central
be accomplished by a

revision of the value for the envelope mass immediately after envelope ejection, and/or by incorporating higher mass loss rates after the superwind phase than assumed in present theoretical models.

The observed positions of the PN in the Hertzsprung-Russell diagram imply that the mass distribution of central stars is less narrow than found earlier and is therefore in better agreement with the mass distribution of white dwarfs. This is also consistent with the observational fact that the local birthrates of PN and white dwarfs are comparable.

The theoretical prediction that high-mass progenitor stars produce high-luminosity PN with higher He and N abundances than average is confirmed by the present observations.

- 6) It is found that for the sample of PN with standard distances the ionized masses vary from $\sim 0.002 M_{\odot}$ to $\gtrsim 0.5 M_{\odot}$. This confirms the results of the study of the PN at the galactic centre.
- 7) The standard distances are used to verify the validity of various statistical distance scales. It is shown that individual distances derived from these scales are generally quite unreliable. This is due to the large spread in intrinsic parameters such as ionized nebular mass and central star luminosity. It is shown that a unique distance scale for all PN cannot be defined. Each of the published scales is in fact a mixture of several scales which represent PN that have originated from stars with different initial masses.